

# FATTY ACIDS CONTENT CONTROL IN FOOD PRODUCTS BY GAS



Institute  
of Animal  
Biology  
NAAS

## CHROMATOGRAPHY

*Liubov Poliuzhyn, Olha Dushna, Olha Korobova*



State Scientific  
Research Control  
Institute of  
Veterinary Medicinal  
Products and Feed  
Additives

State Scientific Research Control Institute of Veterinary Medicinal Products and Feed Additives, Lviv, Ukraine

### Introduction

In different food products there is a certain fatty acid (FA) composition is characterized for. Manufacturers must include the content of fats, proteins, carbohydrates, fats (saturated, monounsaturated and polyunsaturated) and energy value on the labels in accordance with current law. Plant products such as coffee, oils and soy has a specific interrelation of fatty acids. Mainly they are palmitic, stearic, oleic, linoleic and others. For example butter has a specific indicator of fatty acids content such as lauryl, myristic, palmitic, stearic, oleic and linoleic and their ratio, which indicates the quality of the product.

### Material & Methods

The reaction which is the basis of the method:



#### Reagents

Fatty acids methyl esters : lauryl (C12.0), myristic (C14.0), palmitic (C16.0), heptadecanoic (C17.0), stearic (C18.0), oleic (C18.1), linoleic (C18.2), arachidonic (C20.0).

Solvent: hexane

#### Apparatus

Gas chromatograph: HP-6890 Plus;

Detector: flame ionization detector;

Column: ZB-FFAP 30×0,32×0.25

Carrier gas: Helium

Injection volume: 1 µl

Total run time: 10 min

#### Linearity

FAME	Retention time, min	LOD, ppb	LOQ, ppb
C12.0	3,16	0.04	0.11
C14.0	3,68	0.02	0.06
C16.0	4,64	0.04	0.13
C17.0	5,37	0.03	0.09
C18.0	6,36	0.04	0.12
C18.1	6,67	0.04	0.11
C18.2	7,31	0.03	0.08
C20.0	9,45	0.02	0.06

### Results

#### Analysis of butter

FAME	Limits	Sample 1	Sample 2	Sample 3
C16/C12	5,8-14,5	10,54	11,90	10,69
C18/C12	1,9-5,9	2,90	1,64	4,50
C18.1/C14	1,6-3,6	1,82	12,76	2,96
C18.2/C14	0,1-0,5	0,18	5,77	0,30
C18.1+C18.2/C12+ +C14+C16+C18	0,4-0,7	0,39	1,07	0,59




Figure 1

Chromatograms of FAME at different concentrations




Figure 2

Calibration graphs of FAME

### Conclusions

The method was tested during the analysis of food products (coffee, various oils, soy, confectionery, butter). About 20 samples of butter with different fat content were analyzed. The development method was validated by the criteria of specificity, linearity, repeatability and stability.

This method is planned to be used for routine analysis of fatty acids content in food products.